



## Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact [support@jstor.org](mailto:support@jstor.org).

Fig. 1.




Fig. 2.




Fig. 3.



Fig. 4.





Fig. 5.



*Savage sculp.*

II. Benjamini à Broeckhuysen *Med. et Phil. Doct. &c.*  
*OECONOMIA CORPORIS ANIMALIS*,  
 five cogitationes succinctæ de Mente, Corpore, et utriusque conjunctione, juxta Methodum Philosophiæ Cartesianæ, deductæ. *Amstelodami* 1683. 8°.

---

- I. LUMERICUS TERES, *or some Anatomical Observations on the Round Worm bred in human bodies.* By Edward Tyson M. D. Col. Med. Lond. nec non. Reg. Societ. Soc.

HAVING been so large in my former instance, in my *Discourse on the Joynted-worm*, I intend to Contract myself in *this*. Not that our present subject is scanty, or does not afford a sufficient plenty of remarkable observations; But I chose rather to select what most suites our design. For to be exact and nice in all *particulars*, would require a just Treatise, and exceed the bounds I have at present set my self.

I shall therefore here give the *Anatomy* of the *Lumbricus teres*, that common *Round Worm* which Children usually are troubled with: and in this more particularly make my remarks upon the *Organs* of *generation* in both *Sexes*; and herein shew how vastly different they are from those *parts* in the common *Earth Worms*, and it may be, most others. And withall I had designed, together with this, to have given the *Anatomy* of the *Earth Worm*, but since have altered my intentions: and at present shall refer to the account given of it by the famous *Dr. Willis*, reserving my farther observations of it to another opportunity. This sort of *Worm* by *Hippocrates* is named *serpion*; by *Celsus*, *teres*; and is usually about a foot long, or something more, or less; but I have hitherto observed that the *Male* is generally lesser than the *Female*: so that by  
 their

their *bigness* in the same body I have before *dissection* been able to distinguish the *Sex*. They are about the bigness of a Wheat straw, or a Goose quill; their colour White; but being *subjects* so generally known to all, I shall forbear a further description of their *outward parts*; Onely as I remember I did not observe those *feet*, or asperities on the *Annuli*, as in the *Earth Worm*. At both *extremes* they grow narrow. Their *mouth* is compos'd of three Lips as in our *figure*. So the *Leech* hath three Cartilaginous *Teeth* set in a triangle, by which they make the wound in the Skin in Suction. The *Anus* is a transverse slit a little before the extreme point of the *Tail*.

In opening the body I found I cut thorow a large *Muscle* under the Skin: Which *Muscle* in *Earth Worms* I find is *spiral*; as in a good measure is their *motion* likewise; so that by this means, like the *Worm* of an *amger*, they can the better *bore* their passage into the Earth. Their *reptile* motion also may be explained by a Wire wound on a Cylinder; which when slip't off, and one end extended and held fast, will bring the other nearer it. So the *Earth Worm* having shot out or extended it's body, (which is with a wreathing) it takes hold by those small *feet* it hath, and so contracts the hinder part of it's body.

Likewise I observed that dividing this part there issued out a copious *Ichor*; which is naturally discharged by some *Pores* or small *Vents* in the Skin; which in the *Earth Worm* is of great use, by rendering the surface of the body slippery, that so it might the easierly glide into the Earth. And in these other *Worms* of the *Intestines* this humor (as in *Leeches*) makes a covering to the body, which is often cast off, and observed as a *Mucus*, in the Stools of those troubled with them.

In these *Teretes* of *Animal* bodies I never observed those transverse *Diaphragms* which are so numerous in *Earth Worms*, and do intersect or rather so deeply depress the *Intestine*. But the *Cavity* chiefly seems to be fill'd with

the *genital* parts, which I shall now describe: Onely shall first remark that the passage from the *mouth* was some what straightned for a short space, and was distinguished, as in our figure, from the following *Ductus*; which was a strait *Intestine* continued to the end of the body, without any winding or other distinction of a *Stomack* that I could observe.

As to the *Genital parts* of the *Male*. I could here observe a *Penis*, a *Vesicula seminalis*, and a *Testis*: In the *Female* a *Pudendum*, *Vagina Uteri*, *Cornua Uteri*, and *Spermatick Vessels*.

The *Penis* in the *Male* was placed at the *Tail* or opposite Extreme to the *head*; and seemed to be able to exert it self almost the length of a Barley Corn, or proportionably to the length of the *Vagina* in the *Female*.

At the root of the *Penis* was inserted the neck of the *Vesicula Seminalis*, which gradually grew larger as it ascended in the body, and usually did reach almost half way. 'Twas filled and turgid with a *milkie Juice*; Which it received from a *slender Vessel* of the same Colour inserted into it. Which after one turning, was afterwards very much convoluted; and being so, forms that body I call the *Testis*.

Altho' *this part* be so loosely contexed, as even to the naked Eye it appears but as a continued vessel, and may easily be unravelled it's whole length, which I measured was above a Yard: yet I make no difficulty of giving it the name of a *Testis*; since 'tis now sufficiently known, that the *Testes* in more compleat *Animals* are onely a *congeries* of *Vessels*. And a *Rat*, besides this *Worm*, is not the onely Subject wherein I have found them thus loose and easily separable.

In the *Female Worm*, almost about the middle of the body, but more towards the head, I observed an *Orifice* or *Pudendum*, which led into the *Vagina Uteri*; which soon divided into the two *Cornua* which were large, and remarkable

markable. For descending something winding towards the *Tail*, they were then reflected again, and did each of them terminate in slender Vessels, *White*, as they were, but much smaller; and did lye in several convolutions and windings amongst them. These I take for *Spermatick Vessels*. Having taken those *Vessels*, with the *Cornua Uteri* and *Vagina*, out of the body, and laid them on a Paper to dry; I found from each *Cornu*, to the end of the *Spermatick Vessels* which I had preserved, that they measured above four foot.

I opened the *Cornua Uteri* and found them turgid with a *milky Juice*, having placed a little of it upon a small *Microscope*, I plainly perceived 'twas nothing else but an infinite number of small *Eggs*; tho to the naked eye it appeared onely as a fluid body. These *Eggs* when fresh, appeared, as is represented in our fourth *Figure*, covered with abundance of small asperities; but as they grew dry their Surface appeared Smooth.

By comparing that small quantity I did observe, in which I could distinguish so many *Eggs*, with the whole substance contained in both the *Cornua*, I cannot guess there can be so few as 10000 *Eggs* in each *Female Worm*.

How far different *this Worm* is from common *Earth Worms* as to these *parts*, I need onely to refer to a Dr. *Willis's* figures and account of it, to shew. And I am yet to learn what *Worm* out of the body has these *Organs* thus formed. When once there, the Case is plain how they propagate themselves. And <sup>b</sup> *Menjotius*, and all before him, that were of that Opinion, are mistaken; who say that these *Worms* do not generate; nor have any distinction of *Sexes*. <sup>c</sup> *Hippocrates* is express, αἱ μὲν σαρρήλαι τίνεσι. And I think nothing can be plainer then this distinction of *Sexes* in them.

(a) Willis de *Anima Brutor*. cap. 3.

(b) Ant. Menjotius. *dissertat. Patholog.* part. 3. p. m. 512.

(c) Hippocr. *l. 4. de morbis*.

But I find on the other hand, there are *many* who do not onely allow them to generate, but do make them *Viviparous* too.

Thus P. <sup>d</sup> Borellus tells us, *Vermem Crassum ab hominis Corpore eductum, fortèque pedibus exenteratum, non sine admiratione vidi vermiculis innumeris refertum esse.* So <sup>e</sup> Amatus Lusitanus tells much such a Story; that a Girl voiding a large *Worm*, and the father treading on it, *ex eo alij prodierunt Vermes.* And <sup>f</sup> Fælix Platerus gives an observation of a Boy that was Hydropical, and voided all his excrements upwards; who dying in the *Hospital*, and they observing a motion and palpitation in his belly, were afraid to bury him till they had sent for the *Doctor*. He opening him found the *Intestines* in some places swell'd as big as his Thigh, in others so convoluted, intorted, and twisted, that hindred any passage downwards, either of Excrements or Wind; *Sed & vermibus vivis quamplurimis repleta erant, qui rursus alijs minoribus referti.* You may see an Instance likewise *de Vermibus fætis* in Salmuth Cent. 3. Obs. 24. But <sup>g</sup> Dominicus Panarolus is very express; and tells us he observed it thus in two several persons. *In utroque expulsi fuerunt Vermes Colore Carnei, longitudine circa sexdecem digitos, qui prægnantes erant, & ligno collisi cum fuissent, apparuerunt vermes parvi, subtiles, albi, longitudine sex digitorum, prope innumeri, qui tanquam serpentes parvi movebantur.* But whatever is related of *this* nature I cannot but think it is a *mistake*; and that they were imposed upon by the *Genital parts* of this *Worm*; which not warily examined, might easily make them to think they are so many small *Worms*. For they are not *Viviparous* but *Oviparous*, as I have shewn; and their containing so vast a number of *Eggs* in the *Cornua*

(d) P. Borellus *hist. & observ. Cent. 1. Obs. 89.*

(e) Amatus Lusitanus *Cent. 3. curat. 46.*

(f) Fælix Platerus *Obs. lib. 3. p. m. 657.*

(g) Dom. Panarol, *Obs. Med. Prætec. 5. Obs. 15.*

*Uteri*, as I have expressed, does sufficiently account for that prodigious quantity, that are sometimes observed to be bred in *Animal* bodies.

<sup>b</sup> *Panarolus* tells us he once saw the Stomack and Guts stuffed with them so that they ascended up to the Throat. <sup>i</sup> *Baricellus* by the use of *Crude Mercury* brought away from a Patient above a hundred. <sup>k</sup> *Jo. Jadoc. Weckerus* did the like with the use of *Tansy Seed* and Syrup of *Violets*. *Gabucinus* saw voided by Stool 177. <sup>l</sup> *Benivenius* saw voided by a Child 7 years old 152 Worms. And <sup>m</sup> *Jacob Hollerius*, out of *Musa*, gives us an History of a man 82 years old, who voided above 500. And <sup>n</sup> *Petrus Paulus Pereda* saw a Noble-man's Child in a few days void almost a Thousand, and she voided 40 in 4 hours time.

Those *Animal's* are usually the most *Multiparous*, whose young are the most exposed to danger; and were it not so here that the greatest part of the *litter* of this *Worm* is usually carried forth by the *Feces*, it could not be avoided but we should be devoured by an *Enemy* we breed in our own *Bowels*. That caution therefore of <sup>o</sup> *Henr. ab Heers* I think is necessary. To avoid the giving the Powder of these *Worms* for expelling others, since we cannot be secure but that at the same time we may sow the Seed for propagating more.

(b) *Panarol. Pentec. 1. Obs. 41.*

(i) *Baricel. in hortulo genitali.*

(k) *Wecker. de observ. propriis.*

(l) *Beniven. de abditis. cap. 85.*

(m) *Holler. de morb. intern. lib. 1. in Schol. ad cap. 54.*

(n) *Pereda de curand. morb. lib. 1. cap. 5.*

(o) *Henr. ab Heers Obs. med. l. 1. Obs. 9. p. 101.*



THE  
EXPLANATION  
OF THE  
FIGURES.

---

FIGURE I.

**R**epresents the *Male Worm* opened. Where

- a.* Shews the three Lips of the Worm.
- b.* The *Oesophagus*, or Gullet.
- c c c.* The large Intestine.
- d.* The *Penis*.
- e e.* The *Vesicula Seminalis*.
- f.* The *Testis*.

FIGURE II.

Represents the *Female Worm* opened. Where

- a.* Shews the Mouth.
- b.* The Gullet.
- c c c.* The Intestine, or Gut.
- d d d d.* The *Vagina Uteri*.
- e.* The two *Cornua Uteri*.
- fff.* The Spermatick Vessels.
- g.* The *Anus*.

FIG.

FIGURE III.

Represents the Genital parts of the *Female*  
explicated. Where

- a. Shews the *Pudendum* or *Foramen* as it appears on the out  
side of the Skin.
- b. The *Vagina Uteri*.
- c c. The two *Cornua Uteri*.
- d d. The *Spermatick Vessels*.

FIGURE IV.

Represents the *Eggs* of this Worm, as they appear-  
ed being viewed by the *Microscope*.



Y

*A remark-*